# Journal of Surgical Oncology

## Table of Contents

Volume 82, Number 1, January 2003

**Guest Editorial: Impact of the Surgeon on Cancer Management Outcomes**
James T. Helsper 1

**Anal Sphincter Preservation in Locally Advanced Low Rectal Adenocarcinoma After Preoperative Chemoradiation Therapy and Coloanal Anastomosis**
Pedro Luna-Pérez, Saúl Rodríguez-Ramírez, Freddi Hernández-Pacheco, Marcos Gutiérrez de la Barrera, Raúl Fernández, and Sonia Labastida 3
*This study evaluated feasibility, morbidity, and functional results of anal sphincter preservation after preoperative chemoradiation therapy and coloanal anastomosis in patients with rectal adenocarcinoma located 3–6 cm above the anal verge.*

**Role of Metastasectomy in the Management of Thyroid Carcinoma: The NIH Experience**
Ho Pak, Loukas Gourgiotis, Wen-I Chang, Lori C. Guthrie, Monica C. Skarulis, James C. Reynolds, Maria J. Merino, David S. Schrump, Steven K. Libutti, H. Richard Alexander, and Nicholas J. Sarlis 10
*Extensive targeted metastasectomy, used in conjunction with other nonsurgical treatment modalities, can be beneficial to selected patients with disseminated thyroid carcinoma when persistent or recurrent distant disease exists.*

**Prognostic Significance of the Size of Cancer Nests in Metastatic Lymph Nodes in Human Esophageal Cancers**
Takamichi Komori, Yuichiro Doki, Toshiyuki Kabuto, Osamu Ishikawa, Masahiro Hiratsuka, Yo Sasaki, Hiroaki Ohigashi, Kohei Murata, Terumasa Yamada, Isao Miyashiro, Masayuki Mano, Shingo Ishiguro, and Shingi Imaoka 19
*The clinical significance of the size of cancer nests in metastatic lymph nodes was evaluated for 170 node-positive esophageal cancer patients. The size of the largest cancer nest ($N_{max}$) was an independent prognostic factor, as was the number of positive nodes.*

**Lymphocytic Infiltration Surrounding Liver Metastases From Colorectal Cancer**
Keiichi Okano, Takashi Maeba, Akito Moroguchi, Ken Ishimura, Yukihiko Karasawa, Kunihiko Izuishi, Fuminori Goda, Hisashi Usuki, Hisao Wakabayashi, and Hajime Maeta 28
*Lymphocytic infiltration surrounding metastatic liver tumor was graded for evaluation of clinical significance in patients with colorectal liver metastases. Multivariate analysis identified that the extent of lymphocytic infiltration between the metastatic nodule and hepatic parenchyma is closely related to prognosis in patients who underwent hepatic resection for liver metastasis from a colorectal cancer.*

**Prognostic Significance of Biomarkers in Squamous Cell Carcinoma of the Tongue: Multivariate Analysis**
Hemangini H. Vora, Neelam G. Shah, Devendra D. Patel, Trupti I. Trivedi, and Priya R. Chikhlikar 34
*Expression of a panel of biomarkers was studied in anterior tongue tumors from the oral cavity and in posterior tongue tumors from the oropharynx of patients with early- and locally advanced-stage disease, to evaluate their prognostic value.*

**Enhancement of Integrins by Interleukin-1α, and Their Relationship With Metastatic and Invasive Behavior of Human Pancreatic Ductal Adenocarcinoma Cells**
Hirozumi Sawai, Hiromitsu Takeyama, Minoru Yamamoto, Akiyoshi Furuta, Hitoshi Funahashi, Yuji Okada, Mikinori Sato, Moritsugu Tanaka, and Tadao Manabe 51
*To examine the mechanisms of metastasis formation in metastatic human pancreatic cancer, we investigated the enhanced integrin expression, adhesion, and invasiveness into extracellular matrix proteins of human pancreatic ductal adenocarcinoma cells after exposure to IL-1. Our results indicated that $_{61}$-integrin and IL-1RI expression may play important roles in metastasis formation and tumor invasion.*

**Downregulation of Putative Tumor Suppressor Gene TSC-22 in Human Brain Tumors**
Katherina O. Shostak, Vladimir V. Dmitrenko, Oleg M. Garifulin, Vladimir D. Rozumenko, Olexiy V. Khomenko, Yuriy A. Zozulya, Günther Zehetner, and Vadym M. Kavsan 57
*Significantly decreased level of TSC-22 mRNA production in human brain tumors and, as was shown previously, in salivary gland tumors, as well as the antiproliferative role of TSC-22 protein, strongly suggest a tumor suppressor role for TSC-22.*

**Regional Chemotherapy Is Indicated After Surgical Resection of Colorectal Metastases to the Liver: A Debate**
Nikhil I. Khushalani, Brian Patrick McKinley, John F. Gibbs, Cynthia Gail Leichman, and Nicholas J. Petrelli 65

**How I Do It: Localization of Nonpalpable Breast Lesions Using Hook-Wire Combined with Isosulfan Blue Dye**
George C. Zografos, Carmen Doumitriou, Dimitrios Lappas, George Karatzikos, Emi Bami, Costas Fotiadis, Evagelos Menenakos, and George Androulakis 73

**How I Do It: Preoperative Endoscopic Clipping: Localizing Technique of Early Gastric Cancer**
Keun Won Ryu, Jun Ho Lee, Il Ju Choi, and Jae Moon Bae 75

## Volume 82, Number 2, February 2003

**Guest Editorial: Will We Ever Succeed in Resolving the Adjuvant Treatment Dilemma in Rectal Cancer?**
Nicholas J. Petrelli 79

**Association Between Extent of Axillary Lymph Node Dissection and Patient, Tumor, Surgeon, and Hospital Factors in Patients With Early Breast Cancer**
David W. Petrik, David R. McCready, Carol A. Sawka, and Vivek Goel 84
*The extent of axillary lymph node dissection varies and has been found to be associated with survival. In this study, we found that lower patient age and surgery in an academic setting were associated with a greater number of lymph nodes examined.*

**Intradermal Radioisotope Injection Is Superior to Subdermal Injection for the Identification of the Sentinel Node in Breast Cancer Patients**
Kazuyoshi Motomura, Yoshifumi Komoike, Yoshihisa Hasegawa, Tsutomu Kasugai, Hideo Inaji, Shinzaburo Noguchi, and Hiroki Koyama 91
*The intradermal injection of radiocolloids improves the identification rate of sentinel nodes over the subdermal injection in breast cancer patients. Tin radiocolloid, with a large particle size, will easily migrate into the lymphatics by intradermal injection.*

**Commentary**
Omgo E. Nieweg and Emiel J.Th. Rutgers 96

**Significance of Reduction Surgery in Multidisciplinary Treatment of Advanced Hepatocellular Carcinoma With Multiple Intrahepatic Lesions**
Hisao Wakabayashi, Takafumi Ushiyama, Ken Ishimura, Kunihiko Izuishi, Yukihiko Karasawa, Tsutomu Masaki, Seishiro Watanabe, Shigeki Kuriyama, and Hajime Maeta 98
*By comparing the survival rates of patients treated with or without surgery, the significance of, and the indication for, reduction surgery in the multidisciplinary treatment of patients with hepatocellular carcinoma with multiple intrahepatic lesions were examined. When combined with intraoperative adjuvant therapy for remaining satellite tumors, reduction surgery provided survival benefit for selected patient groups.*

**Prognostic Use of Growth Characteristics of Early Gastric Cancer and Expression Patterns of Apoptotic, Cell Proliferation, and Cell Adhesion Proteins**
Wensheng Pan, Hideaki Ishii, Yoshiro Ebihara, and Glenda Gobe 104
*Selection of suitable treatment for early gastric cancers, such as endoscopic mucosal resection or the major surgery with resection of the cancer and lymph nodes, may be assisted by comparing the growth characteristics of the cancer with selected molecular characteristics. The results could be used to predict those cases that have a higher risk of developing secondary metastases.*

**p53 Alteration Is Not an Independent Prognostic Indicator, But Affects the Efficacy of Adjuvant Chemotherapy in Human Pancreatic Cancer**
Ming Dong, Yoshinori Nio, Kunihiro Yamasawa, Tomoko Toga, Lishuang Yue, and Takayuki Harada 111
*A p53 abnormality was not an independent factor for evaluating the prognosis of patients with pancreatic cancer, but it was a beneficial indicator for selecting a reasonable strategy of adjuvant chemotherapy against pancreatic cancer.*

**Establishment of a Nude Mouse Model of Hepatic Metastasis for Evaluation of Targeted Retroviral Gene Delivery**
Maria Stapfer, Jie Hu, Denise Wei, Susan Groshen, and Robert W. Beart Jr. 121
*The goal of this study was to establish a reproducible nude mouse model of liver metastasis and investigate the potential of deploying targeted injectable retroviral vectors for metastatic gastrointestinal cancer.*

**Commentary**
Michael Morse, Bryan Clary, and H. Kim Lyerly 131

**Heterotopic Ossification in Rectal Cancer: Rare Finding With a Novel Proposed Mechanism**
Alan P. Kypson, Emilie Morphew, Relief Jones, Marcia R. Gottfried, and Hilliard F. Seigler 132
*Expression of a bone morphogenic protein, BMP-2, within the tumor cells of a rectal carcinoma is demonstrated for the first time, revealing a potential signaling mechanism for the transformation of mesenchymal cells into bone.*

**Commentary**
Anthony M. Magliocco 137

**How I Do It: Semi-Closed Hyperthermic-Antiblastic Peritoneal Perfusion (HAPP) in the Treatment of Peritoneal Carcinosis**
Michele De Simone, Riccardo Barone, Marco Vaira, Bartolomeo Aghemo, Pierroberto Mioli, Carlo Franco, Stefano Scuderi, Daniela Costamagna, and Marcello Dei Poli 138

## Volume 82, Number 3, March 2003

**Guest Editorial: What Is a False-Negative Result for Sentinel Node Procedures in Breast Cancer?**
Susanne H. Estourgie, Omgo E. Nieweg, Emiel J.T.H. Rutgers, and Bin B.R. Kroon 141

**Previous Wide Local Excision of Primary Melanoma Is Not a Contraindication for Sentinel Lymph Node Biopsy of the Trunk and Extremity**
Wey L. Leong, Danny M. Ghazarian, and David R. McCready 143
*This prospective case-control study provides data regarding the feasibility of sentinel lymph node biopsy (SLNB) in patients who have already had a wide local excision (WLE) of their primary cutaneous melanoma. SLNB in these patients with a previous WLE was compared in a case-control manner with those having a concurrent SLNB and WLE.*

**Postoperative Taste and Smell Deficit After Upper Gastrointestinal Cancer Surgery—An Unreported Complication**
Adrian M. Harris and S. Michael Griffin 147
*Temporary taste and smell deficit after upper GI cancer surgery is reported as a new complication, but the mechanism remains unexplained. It occurs in nearly one-half of all patients and should be highlighted in preoperative consent for these operations.*

**Commentary**
Eduardo J.B. Ramos, Antonio L. Campos, Julio C.U. Coelho, Undurti Das, Michael M. Meguid, and Paul Cunningham 150

**Immunohistochemically Detected Micrometastasis in Lymph Nodes From Superficial Esophageal Squamous Cell Carcinoma**
Tadashi Tanabe, Tadashi Nishimaki, Hidenobu Watanabe, Yoichi Ajioka, Kohei Akazawa, Shintaro Komukai, and Katsuyoshi Hatakeyama 153
*Nodal micrometastasis from superficial esophageal cancer was found in 26.1% of patients with histologically negative nodes and was significantly correlated with intraesophageal multicentric cancer and venous invasion. Yet nodal micrometastasis had no impact on patient survival.*

**Sensitization of Thermotolerant SCK Cells to Hyperthermia and Freezing With Reduction of Intracellular pH: Implications for Cryosurgery**
Abram H. Burgher, David J. Swanlund, Robert J. Griffin, Chang W. Song, John C. Bischof, and Kenneth P. Roberts 160
*Induction of thermal tolerance does not protect SCK cells against subsequent freezing injury. A low pH environment actually sensitizes thermotolerant cells to freeze-induced cell death.*

**Mitochondrial DNA Alterations in Thyroid Cancer**
Betty C. Tong, Patrick K. Ha, Karan Dhir, Mingzhao Xing, William H. Westra, David Sidransky, and Joseph A. Califano 170
*Alterations in the D310 region of the mitochondrial genome of thyroid cancers were analyzed for alterations. The frequency of alteration is lower than that found in other epithelial tumors and shows no relationship to histologic type or grade.*

**Increased Dihydropyrimidine Dehydrogenase Activity in Breast Cancer**
Keisei Anan, Shoshu Mitsuyama, Keiyoshi Tamae, Nobuhiro Suehara, Kazuyoshi Nishihara, Yoshiaki Ogawa, Yuji Abe, Toshimitsu Iwashita, and Satoshi Toyoshima 174
*Dihydropyrimidine dehydrogenase (DPD) activity is substantially upregulated in breast cancer tissue and is higher than that reported previously. Because cancer therapy adjusted according to the level of DPD expression in patients appears to be a promising approach to optimizing 5-FU chemotherapy, the clinical implications of DPD inhibitors in patients being treated for breast cancer with oral fluoropyrimidines should be further investigated.*

**Cytochrome P-450 Expression and Lipid Peroxidation in Gallbladder Cancer**
Manoj Pandey, Lal B. Sharma, and Vijay K. Shukla 180
*The presence of cytochrome P-450 in gallbladder mucosa was studied and its expression correlated with concentration of lipid peroxidation products.*

**Quantitative Expression of Protein Markers of Plasminogen Activation System in Prognosis of Colorectal Cancer**
Da-Qiang Seetoo, Philip J. Crowe, Pamela J. Russell, and Jia-Lin Yang 184
*The prognostic significance of five protein markers of the plasminogen activation system, urokinase-type plasminogen activator (uPA) and its receptor (uPAR), plasminogen, plasminogen activator inhibitor-1 and -2 (PAI-1 and PAI-2), was assessed quantitatively in paraffin-embedded tissue sections of 56 cases of patients with colorectal cancer. uPA and uPAR were found to be independent predictors of liver metastasis, patient overall survival, and cancer-specific survival in this patient series.*

**Combined Loss of Expression of $O^6$-Methylguanine-DNA Methyltransferase and hMLH1 Accelerates Progression of Hepatocellular Carcinoma**
Shiroh Matsukura, Kohji Miyazaki, Hiroyuki Yakushiji, Akiomi Ogawa, Yongxin Chen, and Mutsuo Sekiguchi 194
*Hepatocellular carcinoma lacking both methylguanine methyltransferase (MGMT) and human Mut L homologue 1 (hMLH1) expression correlates with an advanced stage and a poor prognosis. The expression status of both DNA repair proteins is a predictive prognostic factor in patients with hepatocellular carcinoma after surgical resection.*

**Metastatic Potential of Dermatofibrosarcoma Protuberans With Fibrosarcomatous Change**
Rebecca M. Minter, John D. Reith, and Steven N. Hochwald 201
*A rare case of a DFSP-FS metastasis to the head of the pancreas without any preceding local recurrence is reported.*

**Current Management of Melanoma: Benefits of Surgical Staging and Adjuvant Therapy**
Kelly M. McMasters and Susan M. Swetter 209
*Sentinel lymph node biopsy is appropriate for pathologic nodal staging of many patients with melanoma. Node-positive patients benefit from therapeutic lymphadenectomy, and from adjuvant therapy with high-dose interferon alfa-2b.*

## Volume 82, Number 4, April 2003

**Editorial Commentary: Molecular Staging of Cancer: Concepts of Today, Therapies of Tomorrow**
Heike Allgayer 217

**Detection of Disseminated Tumour Cells in Bone Marrow of Patients With Isolated Liver Metastases From Colorectal Cancer**
Kristin Bjørnland, Kjersti Flatmark, Tom Mala, Øystein Mathisen, Arne Bakka, Ansgar O. Aasen, Anstein Bergan, Odd Søreide, and Øystein Fodstad 224
*Bone marrow from patients with isolated liver metastases from colorectal cancer was examined for micrometastases using immunomagnetic beads. The frequency of bone marrow micrometastases was low (8%), and bone marrow status should not affect current treatment protocols.*

**Surgery-Related Shedding of Breast Cancer Cells as Determined by RT-PCR Assay**
Xi-Chun Hu, Wings T.Y. Loo, and Louis W.C. Chow 228
*Breast cancer operations may result in the shedding of cancer cells into the circulation. However, these cells may not be involved in the formation of metastasis.*

**Commentary**
Cary S. Kaufman 233

**Impact of Adjuvant Chemotherapy in the Long-Term Outcome of Patients With Resected Gastric Cancer**
Juan J. Grau, Marta Martín, José Fuster, Manuel Pera, Juan C. García-Valdecasas, Josep A. Bombí, José M. Bordas, Felipe Alcobendas, Luis Grande, and Jordi Estapé 234
*Survival results of adjuvant postoperative chemotherapy with mitomycin C alone or combined with Tegafur in 314 resected gastric cancer patients are reported. The two drugs are more effective than one drug regimen, as confirmed by multivariate analysis.*

**Metastatic Pattern of Lymph Node and Surgery for Gastric Stump Cancer**
Shao-Liang Han, Ya-Wei Hua, Cheng-Hu Wang, She-Qing Ji, and Jing Zhuang 241
*The present study showed that the pattern of lymph node (LN) spread in gastric stump cancer (GSC) was different from carcinoma of the "primary" stomach (including tumor of the upper third). Our results imply that gastric stump cancer has a wide range of LN metastases, including LN within jejunal mesentery in B-II reconstruction cases, and curable resection may obtain better results. Therefore, we suggest that radical operation for B-I patients needs removal of gastroduodenectomy anastomosis and the above LNs, and that B-II patients need removal of 10 cm of jejunum beside gastrojejunostomy anastomosis, and clearance of LN within its mesentery, in addition to B-I GSC.*

**Prognostic Impact of DNA Ploidy Pattern, S-Phase Fraction (SPF), and Proliferating Cell Nuclear Antigen (PCNA) in Patients With Primary Gastric Lymphoma**
Chrysoula J. Belessi, Aikaterini S. Parasi, Helen S. Manioudaki, Nikolaos P. Laoutaris, Nikolaos C. Legakis, Georgios Th. Peros, and Georgios A. Androulakis 247
*S-phase fraction and proliferative cell nuclear antigen are important prognostic factors in patients with primary gastric lymphomas. DNA ploidy had no impact on survival in patients with primary gastric lymphomas in the present study.*

**Role of Skip Metastasis to Mediastinal Lymph Nodes in Non-Small Cell Lung Cancer**
Klaus L. Prenzel, Stefan P. Mönig, Jan M. Sinning, Stefan E. Baldus, Christian A. Gutschow, Guido Grass, Paul M. Schneider, and Arnulf H. Hölscher 256
*In a retrospective study the role of skip metastasis in NSCLC was analyzed. Patients with skip metastasis showed a significantly longer survival than those with continuous metastatic spread.*

**Cardiac Injury During Needle Localized Surgical Breast Biopsy**
Steve R. Martinez, Mark Gelfand, Hisham S. Hourani, Joseph J. Sorrento, and Eugene P. Mohan 261
*Needle localized breast biopsy is commonly used to diagnose and remove non-palpable breast lesions. We present the first case of cardiac injury resulting from needle localized breast biopsy.*

**Abstracts From the 3rd World Congress of the World Federation of Surgical Oncology Societies, Los Angeles, California, USA March 6–9, 2003** 266

**Author Index to Volume 82** 281

**Subject Index to Volume 82** 283